

HOMOCLINIC SOLUTIONS FOR FOURTH ORDER TRAVELING WAVE EQUATIONS

SANJIBAN SANTRA, JUNCHENG WEI

ABSTRACT. We consider homoclinic solutions of fourth order equations

$$u'''' + \beta^2 u'' + V_u(u) = 0 \text{ in } \mathbb{R},$$

where $V(u)$ is either the suspension bridge type $V(u) = e^u - 1 - u$ or Swift-Hohenberg type $V(u) = \frac{1}{4}(u^2 - 1)^2$. For the suspension bridge type equation, we prove existence of a homoclinic solution for *all* $\beta \in (0, \beta_*)$ where $\beta_* = 0.7427 \dots$. For the Swift-Hohenberg type equation, we prove existence of a homoclinic solution for each $\beta \in (0, \beta_*)$, where $\beta_* = 0.9342 \dots$. This partially solves a conjecture of Chen–McKenna [12].

1. INTRODUCTION

The study of homoclinic and heteroclinic solutions for fourth order equations has attracted a lot of attention for the last two decades. Though simple-looking, the fourth order equations appear to be difficult and pose lots of very challenging questions. We refer to the survey papers [21] and the monograph [24] for further references.

Motivated by the appearance of traveling wave behavior on the Narrows Tacoma bridge and the Golden Gate bridge, McKenna and Walter [22] considered the following nonlinear beam equation

$$w_{tt} + w_{xxxx} + V_w(w) = 0$$

where V_w is the restoring force and is chosen such that the effective force of the cables holds the beam up but the constant force of gravity holds it down on the assumption that there is no reaction force due to compression. Here $w(x, t)$ denotes the displacement of the beam from the unloaded state [19]. This leads to a fourth order beam equation

$$(1.1) \quad \frac{\partial^2 w}{\partial t^2} + \frac{\partial^4 w}{\partial x^4} = -w^+ + 1 \text{ in } \mathbb{R}$$

where $w^+ = \max\{w, 0\}$. Note that (1.1) also arise in the study of deflection in railway tracks and undersea pipelines. See [1] and [8].

If we look for a traveling wave solution of the type $w(x, t) = 1 + u(x - \beta t)$, then (1.1) is transformed to a fourth order differential equation of the form

$$(1.2) \quad u'''' + \beta^2 u'' + (u + 1)^+ - 1 = 0 \text{ in } \mathbb{R}$$

1991 *Mathematics Subject Classification.* Primary 34B15, 34B60, 34E18.

Key words and phrases. Bistable nonlinearity, mountain pass solution, Morse index, homoclinic solutions, solitons.

where β denotes the wave speed. McKenna–Walter [22] studied (1.2) by solving an ordinary differential equation explicitly as

$$(1.3) \quad \begin{cases} u'''' + \beta^2 u'' + u = 0 & \text{if } u \geq -1, \\ u'''' + \beta^2 u'' = 1 & \text{if } u \leq -1 \end{cases}$$

and then glued the two solutions to match at $u = -1$ (called one-trough solutions). In fact, they noticed that as the wave speed approaches $\sqrt{2}$, the solution becomes highly oscillatory in nature, and as β approaches 0, they appear to go to infinity in amplitude. It was also noticed by numerical experiments that some of the traveling wave solutions appear to be stable, that is, when two waves collide, they pass through each other like solitons having many nodes.

Later on, Chen–McKenna [11] applied mountain pass theorem on $H^2(\mathbb{R})$ to prove that (1.2) has a nontrivial solution. In addition the calculations in [22] suggest that there are many solutions, possibly infinitely many solutions, though it is only known that there exists at least one non-trivial solution. In [9], Champneys and McKenna proved that there exist $0 < \beta' < \beta'' < \sqrt{2}$ such that (1.2) has infinitely many *multitroughed* homoclinic solutions for all $\beta \in (\beta', \beta'')$ using the ideas in [7], [14] and [27].

The model (1.2) has some serious drawbacks. Firstly, it simplifies the nonlinearity of the physical situation, by not allowing nonlinear effects until the deflection is quite large. Secondly, the non-smoothness of the restoring force leads to numerical difficulties. So the following modified version of (1.2) was proposed in [11]

$$(1.4) \quad \begin{cases} u'''' + \beta^2 u'' + e^u - 1 = 0 & \text{in } \mathbb{R}, \\ u \neq 0 \\ u \in H^2(\mathbb{R}). \end{cases}$$

Though the nonlinearity in (1.4) looks similar to that in (1.2), the study of (1.4) is quite difficult. In addition, $V(u) = \int_0^u (e^t - 1)dt = e^u - u - 1$ is not symmetric and it has linear growth at $-\infty$ and grows like e^u at $+\infty$.

In [29], Smets–van den Berg used mountain-pass lemma and Struwe’s monotonicity trick [30] to prove that for *almost all* $\beta \in (0, \sqrt{2})$, (1.4) admits a solution. Later on in [5], Breuer, Horák, McKenna and Plum used a computer assisted proof to conclude that if $\beta = 1.3$, there is at least 36 solutions. It was then conjectured in [5] that there is at least one homoclinic solution for *all* $\beta \in (0, \sqrt{2})$. In this paper, we partially solve this conjecture.

Theorem 1.1. *There exists $0 < \beta_\star < 1$ such that for all $\beta \in (0, \beta_\star)$, (1.4) admits a homoclinic solution and u decays in the form $e^{-\tau(\beta)|x|} \cos(ax + b)$ for some $a, b \in \mathbb{R}$ and $\tau(\beta) > 0$. (Explicitly, $\beta_\star \sim 0.7427 \dots$).*

We will also consider the Swift–Hohenberg equation which is a general model for pattern-forming process derived in [31] to describe random thermal fluctuations in the Boussinesque equation and in the propagation of lasers [20]. It also arises in the study of ternary mixtures made up of oil, water and surfactant agents yielding a free energy functional of the Ginzburg–Landau equation given by,

$$(1.5) \quad \Psi(u) = \int_{\mathbb{R}^3} [(\Delta u)^2 + h(u)|\nabla u|^2 + V(u)]dx$$

where the scalar parameter u is related to the local difference of the concentration of oil and water [18]. The function h denotes the amphilic properties and $V(u)$ denotes the potential (the bulk free energy of the ternary mixture) [2]. Not only they have important applications in science especially in statistical mechanics of self avoiding surfaces, but also in cell membrane biology, in string theory and in high energy physics [28]. The existence of heteroclinic solution has been studied extensively in [4] when h changes sign.

In this paper we also consider

$$(1.6) \quad u'''' + \beta^2 u'' + V_u(u) = 0 \text{ in } \mathbb{R}$$

where $V_u(u) = -u + u^3$. For this model, a question of interest is phase transition i.e. solutions connecting to $u = \pm 1$. Peletier and Troy studied homoclinic and heteroclinic solutions when $h(u) = -\beta^2$ in [25], [26], though nothing is known about the existence of heteroclinic solutions of (1.6) for $0 < \beta < \sqrt{8}$. Buffoni [6] proved that if $V_u(u) = -u + u^2$, then (1.6) admits at least one solution for all $\beta \in (0, \sqrt{2})$.

Our techniques in proving Theorem 1.1 actually allows to conclude similar results for the well-known Swift-Hohenberg model

$$(1.7) \quad \begin{cases} u'''' + \beta^2 u'' + u(u^2 - 1) = 0 & \text{in } \mathbb{R} \\ u - 1 \in H^2(\mathbb{R}). \end{cases}$$

Smets-van den Berg [29] proved that for *almost all* $\beta \in (0, \sqrt{8})$, problem (1.7) has a homoclinic solution. For (1.7), we have

Theorem 1.2. *For each $\beta \in (0, \beta_0)$, where $\beta_0 \approx 0.9342 \dots$, (1.7) admits a homoclinic solution.*

Here $\beta_0 = \sqrt{\frac{\sqrt{2}}{k_0}}$ where $4k_0^2 - 2k_0 - 3 = 0$. In particular, $\beta_0 \approx 0.9342 \dots$. As far as we know, Theorems 1.1 and 1.2 are the first result in establishing the existence of homoclinic solutions for explicit β 's.

Let us recall some of the difficulties associated to problem (1.4):

(a) The functional associated to (1.4) does not satisfy the global *Ambrosetti-Rabinowitz condition*, i.e.

$$V_t(t)t - \theta V(t) \geq 0$$

for some $\theta > 2$ and for all $t \in \mathbb{R}$. This poses a major problem in proving the boundedness of a Palais-Smale sequence in the H^2 -norm. In [29], Struwe's monotonicity trick was used to conclude the H^2 -boundedness.

(b) Linearizing the equation (1.4) at $u = 0$ we obtain

$$(1.8) \quad w'''' + \beta^2 w'' + w = 0.$$

The roots of (1.8) are given by

$$(1.9) \quad \mu_{\pm}^2 = \frac{-\beta^2 \pm \sqrt{\beta^4 - 4}}{2}.$$

Note that if $\beta \geq \sqrt{2}$, then $\mu_{\pm}^2$ are real and (1.4) can be written as

$$u'''' + \beta^2 u'' + u + e^u - u - 1 = 0$$

and hence can be decomposed into a system

$$(1.10) \quad \begin{cases} u'' - \mu_+^2 u = w & \text{in } \mathbb{R} \\ w'' - \mu_-^2 w = 1 - u - e^u & \text{in } \mathbb{R}. \end{cases}$$

This formulation in fact helps us to obtain a-priori estimates for u and w using strong maximum principle. But if $0 < \beta < \sqrt{2}$ we cannot apply this method to reduce to systems, and in fact monotone homoclinics cannot exist in this range.

(c) Let $w = u''$. Then (1.4) can be written as $w'' + cw = 1 - e^u$ where $c = \beta^2 > 0$. As a result, we cannot apply maximum principle and we cannot say whether a solution of (1.4) after a certain stage is positive or negative.

(d) A solution of (1.4) tends to oscillate infinitely many times even if we have a bound on the Morse index of the solution. This poses a lot of trouble in obtaining solutions converging to zero as $x \rightarrow \pm\infty$.

Our main idea of proving Theorem 1.1 is to bound the H^2 norm by the energy and the *Morse index*. A crucial tool is the Morse index of the mountain-pass solutions. We believe that a more refined analysis should cover the full range $\beta \in (0, \sqrt{2})$.

Finally let us mention that the idea of using Morse index to bound solutions has been used in several recent papers for second order elliptic equations. See Dancer [13] and Farina [16] and the references therein. However, this idea has never been used in fourth order equations because the Moser iteration does not work.

Notations: Throughout this paper, by the equality $B = O(A)$ we mean that there exists $C > 0$ such that $|B| \leq CA$ and $C \approx A$ means that $C = A + o(1)$ where $o(1) \rightarrow 0$ as $\beta \rightarrow 0$.

2. PRELIMINARIES

In this section, we prove existence of a mountain-pass solution and show that its Morse index is *at most one*. This will be used crucially in the next section. We first recall the following definition.

Definition 2.0.1. Let H be a Hilbert space and B be a closed set of H . Let $\mathcal{F}$ be a family of compact subsets of H . Then we call $\mathcal{F}$ a *homotopy stable family with boundary B* if

- (a) Every $A \in \mathcal{F}$ contains B .
- (b) For any $A \in \mathcal{F}$ and any $\eta \in C(H \times [0, 1]; H)$ with $\eta(x, t) = x$ and for all $(x, t) \in (H \times \{0\}) \cup (B \times [0, 1])$ implies that $\eta(A \times [0, 1]) \in \mathcal{F}$.

Definition 2.0.2. A family $\mathcal{F}$ of G -subsets is said to be G -homotopic of dimension N with boundary B if there exists a compact G -subset D of $\mathbb{R}^N$ containing a closed subset D_0 and a continuous G -invariant map $\sigma' : D \rightarrow B$ such that

$$\mathcal{F} = \{A \subset H : A = f(D) \text{ for some } f \in C_G(D, H) \text{ with } f = \sigma' \text{ on } D_0\}.$$

Define

$$K_c = \{u \in H : I(u) = c ; \langle I'(u), u \rangle = 0\}.$$

A Lie group G is said to be a free action if $gx = x$ implies $g = i_d$ for any $x \in H$. We borrow the following lemma from [15] on page 232.

Lemma 2.1. *Let G be a compact Lie group acting freely and differentiably on H . Let I be a G -invariant functional on H and $\mathcal{F}$ be a G -homotopic of dimension N stable with boundary B . If I satisfies $(PS)_c$ where $c := c(I, \mathcal{F})$ and $I''(u)$ is a Fredholm for each level c and $\sup_B I < c$. Then there exists $u \in K_c$ with Morse index of u at most N .*

Proof. For the proof, see [17], Chapter 10. $\square$

We define

$$(2.1) \quad I_\beta(u) = \frac{1}{2} \int_{\mathbb{R}} |u''|^2 dx - \frac{\beta^2}{2} \int_{\mathbb{R}} |u'|^2 dx + \int_{\mathbb{R}} (e^u - u - 1) dx, \quad \forall u \in H^2(\mathbb{R}).$$

First note that I_β is $C^2(H^2(\mathbb{R}))$ and it does not satisfy Palais-Smale condition due to translation invariance of the functional. Moreover, if u is a critical point of I_β , then u is a classical solution of (1.4). Also we have $I_\beta(0) = 0$.

Lemma 2.2. *There exist $r > 0, c > 0$ such that $I_\beta(u) \geq c\|u\|_{H^2(\mathbb{R})}^2$ for all $u \in B_r(0)$ where B_r is a ball centered at the origin in $H^2(\mathbb{R})$. In fact, we can choose r and c to be independent of β .*

Proof. If $\|u\|_{H^2(\mathbb{R})} < r$, then by Sobolev embedding theorem $\|u\|_{L^\infty(\mathbb{R})} < Cr$ for some $C > 0$ (independent of r). Let us choose r so small that if $\|u\|_{H^2(\mathbb{R})} < r$, then $(e^u - u - 1) \geq (\frac{1}{2} - \eta)u^2$ for some $\eta = \frac{2-\beta^2}{12}$. Let $\hat{u}(\xi)$ be a Fourier transform of $u(x)$. Taking Fourier transform we have

$$\begin{aligned} I_\beta(u) &\geq \frac{1}{2} \int_{\mathbb{R}} (\xi^4 - \beta^2 \xi^2 + 1) (\hat{u}(\xi))^2 d\xi - \eta \int_{\mathbb{R}} \hat{u}^2(\xi) d\xi \\ &\geq \frac{1}{2} \int_{\mathbb{R}} (\xi^4 + \xi^2 + 1 - (\beta^2 + 1)\xi^2) (\hat{u}(\xi))^2 d\xi - \eta \int_{\mathbb{R}} \hat{u}^2(\xi) d\xi \\ &\geq \frac{1}{2} \int_{\mathbb{R}} \left(\xi^4 + \xi^2 + 1 - \frac{(\beta^2 + 1)}{3} (\xi^4 + \xi^2 + 1) \right) (\hat{u}(\xi))^2 d\xi \\ &\quad - \eta \int_{\mathbb{R}} \hat{u}^2(\xi) d\xi \\ &\geq \frac{2 - \beta^2}{6} \int_{\mathbb{R}} (\xi^4 + \xi^2 + 1) (\hat{u}(\xi))^2 d\xi - \eta \|u\|_{H^2(\mathbb{R})}^2 \\ (2.2) \quad &= \frac{2 - \beta^2 - 6\eta}{6} \|u\|_{H^2(\mathbb{R})}^2 = \frac{2 - \beta^2}{12} \|u\|_{H^2(\mathbb{R})}^2. \end{aligned}$$

$\square$

Lemma 2.3. *There exists e (independent of β) $\in H^2(\mathbb{R})$ such that $I_\beta(e) < 0$.*

Proof. Choose $v \in H^2(\mathbb{R})$ such that v has compact support and $v < 0$. Let $u_\lambda(x) = v(\lambda x)$, $\lambda > 0$. Then it is possible to choose a $\lambda > 0$ such that

$$\int_{\mathbb{R}} |u_\lambda''|^2 - \beta^2 |u_\lambda'|^2 = -\delta < 0.$$

For $\{u_\lambda < 0\}$ we have $e^{u_\lambda} - u_\lambda - 1 > 0$. Now consider

$$\begin{aligned} \frac{I_\beta(tu_\lambda)}{t^2} &= \frac{1}{2} \int_{\mathbb{R}} (|u_\lambda''|^2 - \beta^2 |u_\lambda'|^2) + \int_{u_\lambda < 0} \frac{e^{tu_\lambda} - tu_\lambda - 1}{t^2} dx \\ (2.3) \quad &= -\frac{\delta}{2} + \int_{u_\lambda < 0} \frac{e^{tu_\lambda} - tu_\lambda - 1}{t^2} dx. \end{aligned}$$

But note that the second term is an integral over a bounded domain as support of u is compact and $\frac{e^{tu_\lambda} - tu_\lambda - 1}{t^2} \rightarrow 0$ as $t \rightarrow \infty$. This implies that

$$I_\beta(tu_\lambda) \rightarrow -\infty \text{ as } t \rightarrow +\infty.$$

Hence the result follows. $\square$

Choose $e = tu_\lambda$. From [29] we know that for almost all $\beta \in (0, \sqrt{2})$, I_β satisfies Palais-Smale condition and hence there exists a mountain pass critical value c_β and

$$c_\beta = \inf_{\gamma \in \Gamma} \max_{t \in [0,1]} I_\beta(\gamma(t)) > 0$$

where

$$\Gamma = \{\gamma : \gamma \in C([0,1], H^2(\mathbb{R})); \gamma(0) = 0, \gamma(1) = e\}$$

and $I_\beta(e) < 0$.

Lemma 2.4. *There exists a constant $C > 0$ independent of β such that $c_\beta \leq C$.*

Proof. We have $I_\beta(e) < 0$. Define a path $\gamma : [0,1] \rightarrow H^2(\mathbb{R})$ such that $\gamma_1(\bar{t}) = \bar{t}e$. Then

$$c_\beta \leq \max_{\bar{t} \in [0,1]} I_\beta(\gamma_1(\bar{t})) \leq C.$$

Hence c_β is uniformly bounded. $\square$

Remark 2.5. Also $I_\beta''(u)$ can be expressed as $I_d - K$ where I_d is the identity map and K is a compact operator. Let $G = \{i_d\}$; the trivial group consisting of the identity element. I_β is a G -invariant functional which satisfies Palais Smale condition [29], for almost all $\beta \in (0, \sqrt{2})$. Choose $B = \{0, e\}$ and let $\mathcal{F}_0^e$ be the collection of all paths joining 0 and e . Then $\mathcal{F}_0^e$ is a homotopy stable family with boundary B . Moreover, $\sup_B I_\beta < c_\beta$. Hence by Lemma 2.1, the solution u_β found in [29] has Morse index at most one. Also note that c_β is a decreasing function of β .

We summarize the results in the following theorem

Theorem 2.6. *For almost all $\beta \in (0, \sqrt{2})$, there exists a mountain-pass solution u_β of (1.4) such that*

- (1) $0 < c_\beta = I_\beta(u_\beta) < C$, where C is independent of $\beta \in (0, \sqrt{2})$,
- (2) u_β has Morse index at most one and $u_\beta \in H^2(\mathbb{R})$,
- (3) the following pointwise identity holds

$$(2.4) \quad u'(x)u'''(x) - \frac{(u''(x))^2}{2} + \frac{\beta^2}{2}(u'(x))^2 + e^{u(x)} - u(x) - 1 = 0.$$

We will call (2.4) a kind of Pohozaev identity which follows by multiplying (1.4) with u' and then integrating in $(-\infty, x)$.

3. KEY INEQUALITIES

In this section, we prove the following key inequalities which will be used to bound the part where u is large.

Lemma 3.1. *Let $k_1 > 1$ be such that*

$$(3.1) \quad k_1^2 - 1 - k_1 - \sqrt{k_1^2 - 1} = 0.$$

Then we have

$$(3.2) \quad \int_{-a}^a (u'')^2 - \beta^2 \int_{-a}^a (u')^2 + \frac{k_1^2 \beta^4}{4} \int_{-a}^a u^2 \geq 0$$

for all $u \in H^2(-a, a)$ and $u(-a) = u(a)$.

Proof. First, we note that the following inequality holds

$$(3.3) \quad \int_{-a}^a (u'')^2 - \beta^2 \int_{-a}^a (u')^2 + \frac{k^2 \beta^4}{4} \int_{-a}^a u^2 \geq 0$$

for all $k \geq 1, u \in H^2(-a, a), u(-a) = u(a) = 0$. See Lemma 5 of [3].

Hence we may assume that $u(-a) = u(a) \neq 0$. Then by rescaling, we may assume that $\beta^2 = 2$ and $u(\pm a) = 1$. We consider the following minimization problem

$$(3.4) \quad M_a = \min_{u \in H^2(-a, a), u(\pm a) = 1} \int_{-a}^a (u'')^2 - 2 \int_{-a}^a (u')^2 + k^2 \int_{-a}^a u^2.$$

Using the inequality (3.3), it is easy to see that the minimizer in (3.4) exists and satisfies

$$(3.5) \quad \begin{cases} u'''' + 2u'' + k^2 u = 0 & \text{in } (-a, a), \\ u(\pm a) = 1, u''(\pm a) = 0. \end{cases}$$

We can assume that u is even since by (3.3), the solution to

$$(3.6) \quad \begin{cases} u'''' + 2u'' + k^2 u = 0 & \text{in } (-a, a) \\ u(\pm a) = 0, u''(\pm a) = 0 \end{cases}$$

is zero, if $k > 1$. From (3.5), we conclude

$$(3.7) \quad u(x) = A \cosh \lambda x \cos \mu x + B \sinh \lambda x \sin \mu x$$

where $r = \lambda + i\mu$ are the roots of $r^4 + 2r^2 + k^2 = 0$. Then the minimum can be computed

$$(3.8) \quad M_a = 2 \left[\int_0^a (u'')^2 - 2 \int_0^a (u')^2 + k^2 \int_0^a u^2 \right] = -2u(a)(u'''(a) + 2u'(a)).$$

In order to show that $M_a \geq 0$ we proceed to calculate A, B .

First we compute the derivatives of u ;

$$u'(x) = (\lambda A + \mu B) \sinh \lambda x \cos \mu x + (B\lambda - \mu A) \cosh \lambda x \sin \mu x,$$

$$u''(x) = ((\lambda^2 - \mu^2)A + 2\lambda\mu B) \cosh \lambda x \cos \mu x + (B(\lambda^2 - \mu^2) - 2\mu\lambda A) \sinh \lambda x \sin \mu x,$$

$$\begin{aligned} u'''(x) &= (\lambda(\lambda^2 - \mu^2)A + 2\lambda^2\mu B + \mu B(\lambda^2 - \mu^2) - 2A\lambda\mu^2) \sinh \lambda x \cos \mu x \\ &\quad + (\lambda B(\lambda^2 - \mu^2) - 2A\lambda^2\mu - 2\mu(\lambda^2 - \mu^2)A - 2B\lambda\mu^2) \cosh \lambda x \sin \mu x. \end{aligned}$$

Using $u''(a) = 0$ we have

$$(3.9) \quad \begin{aligned} u''(a) &= ((\lambda^2 - \mu^2)A + 2\lambda\mu B) \cosh \lambda a \cos \mu a \\ &\quad + (B(\lambda^2 - \mu^2) - 2\mu\lambda A) \sinh \lambda a \sin \mu a \end{aligned}$$

and from $u(a) = 1$ we get

$$(3.10) \quad \begin{cases} A \cosh \lambda a \cos \mu a + B \sinh \lambda a \sin \mu a = 1 \\ 2B\lambda\mu \cosh \lambda a \cos \mu a - 2A\lambda\mu \sinh \lambda a \sin \mu a = (\mu^2 - \lambda^2) \end{cases}$$

which implies that

$$(3.11) \quad A \cosh \lambda a \cos \mu a + B \sinh \lambda a \sin \mu a = 1$$

$$(3.12) \quad -A \sinh \lambda a \sin \mu a + B \cosh \lambda a \cos \mu a = \frac{\mu^2 - \lambda^2}{2\lambda\mu}.$$

By simple computations we obtain

$$A = \frac{\cosh \lambda a \cos \mu a - \frac{\mu^2 - \lambda^2}{2\lambda\mu} \sinh \lambda a \sin \mu a}{\cosh^2 \lambda a \cos^2 \mu a + \sinh^2 \lambda a \sin^2 \mu a}$$

and

$$B = \frac{\sinh \lambda a \sin \mu a + \frac{\mu^2 - \lambda^2}{2\lambda\mu} \cosh \lambda a \cos \mu a}{\cosh^2 \lambda a \cos^2 \mu a + \sinh^2 \lambda a \sin^2 \mu a}.$$

Now

$$(3.13) \quad \begin{aligned} u'''(a) + 2u'(a) &= A(\lambda(\lambda^2 - \mu^2) - 2\lambda\mu^2 + 2\lambda) \sinh \lambda a \cos \mu a \\ &+ B(\mu(\lambda^2 - \mu^2) + 2\lambda^2\mu + 2\mu) \sinh \lambda a \cos \mu a \\ &+ A(-2\lambda^2\mu - \mu(\lambda^2 - \mu^2) - 2\mu) \cosh \lambda a \sin \mu a \\ &+ B(-2\lambda\mu^2 + \lambda(\lambda^2 - \mu^2) + 2\lambda) \cosh \lambda a \sin \mu a. \end{aligned}$$

As a result we have

$$(3.14) \quad \begin{aligned} u'''(a) + 2u'(a) &= \frac{1}{4\lambda} [2\lambda^2(\lambda^2 - \mu^2) - 4\lambda^2\mu^2 + 4\lambda^2 + (\lambda^2 - \mu^2 + 2\lambda\mu + 2)(\mu^2 - \lambda^2)] \sinh 2\lambda a \\ &+ \frac{1}{4\mu} [(\lambda^2 - \mu^2 - 2\lambda\mu + 2)(\mu^2 - \lambda^2) - 2\mu(2\lambda^2\mu + \mu(\lambda^2 - \mu^2) + 2\mu)] \sin 2\mu a \\ &= \frac{1}{4\lambda} [(\lambda^2 + \mu^2)(\lambda^2 - \mu^2) - 4\lambda^2\mu^2 + 4\lambda^2 + (2\mu\lambda + 2)(\mu^2 - \lambda^2)] \sinh 2\lambda a \\ &- \frac{1}{4\mu} [(\lambda^2 + \mu^2)(\lambda^2 - \mu^2) - 4\lambda^2\mu^2 + 4\mu^2 + (2\mu\lambda - 2)(\mu^2 - \lambda^2)] \sin 2\mu a. \end{aligned}$$

Since $r = \lambda + i\mu$ is a root of $r^4 + 2r^2 + k^2 = 0$, we have

$$(3.15) \quad (\lambda^2 - \mu^2)^2 - 4\lambda^2\mu^2 + 2(\lambda^2 - \mu^2) + k^2 = 0.$$

Let $(\lambda^2 - \mu^2) = -1$. Then from (3.15) we have $4\lambda^2\mu^2 = k^2 - 1$ and hence $(\lambda^2 + \mu^2)^2 = k^2$. Hence from (3.13) we have

$$(3.16) \quad -(u'''(a) + 2u'(a)) = -\frac{1}{4}(k - k^2 + 1 + \sqrt{k^2 - 1}) \left[\frac{1}{\lambda} \sinh 2\lambda a - \frac{1}{\mu} \sin 2\mu a \right].$$

Now we determine the sign of $(\sinh 2\lambda a - \frac{\lambda}{\mu} \sin 2\mu a)$. Note that we have $\lambda^2 + \mu^2 = k$ and $\lambda^2 - \mu^2 = -1$. Hence we have $\mu = \sqrt{\frac{k+1}{2}}$ and $\lambda = \sqrt{\frac{k-1}{2}}$.

Let $x = 2\mu a$. Then we have

$$\sinh 2\lambda a - \frac{\lambda}{\mu} \sin 2\mu a = \sinh \frac{\lambda}{\mu} x - \frac{\lambda}{\mu} \sin x.$$

But we know that

$$\sinh \frac{\lambda}{\mu} x > \frac{\lambda}{\mu} x > \frac{\lambda}{\mu} \sin x \quad \forall x.$$

Hence

$$M_a = -2u(a)(u'''(a) + 2u'(a)) \geq 0$$

provided $k^2 - k - 1 - \sqrt{k^2 - 1} \geq 0, k > 1$. This proves (3.2). $\square$

Lemma 3.2. *Let k_2 be such that*

$$(3.17) \quad 4k_2^2 - 2k_2 - 3 = 0, k_2 > 1.$$

Then we have

$$(3.18) \quad \int_a^\infty (u'')^2 - \beta^2 \int_{-a}^\infty (u')^2 + \frac{k_2^2 \beta^4}{4} \int_a^\infty u^2 \geq 0$$

for all $u \in H^2(a, +\infty)$.

Proof. As before, we may assume that $a = 0, u(0) = 1, \beta^2 = 2$. Using the inequality (3.3), it is easy to see that the minimizer for the problem

$$(3.19) \quad M'_a = \min_{u \in H^2(0, \infty), u(0)=1} \int_0^\infty (u'')^2 - 2 \int_0^\infty (u')^2 + k^2 \int_0^\infty u^2$$

exists and satisfies

$$(3.20) \quad \begin{cases} u'''' + 2u'' + k^2 u = 0 & \text{in } (0, +\infty) \\ u(0) = 1, u''(0) = 0. \end{cases}$$

Hence

$$(3.21) \quad u(x) = Ae^{-\lambda x} \cos \mu x + Be^{-\lambda x} \sin \mu x$$

where $r = -\lambda + i\mu$ are the roots of $r^4 + 2r^2 + k^2 = 0$. Similar computations as in Lemma 3.1 give

$$u'''(0) + 2u'(0) \geq 0$$

if $4k^2 - 2k - 3 \geq 0, k > 1$. $\square$

Remark 3.3. Similar results as Lemma 3.2 holds for $u \in H^2(-\infty, -a)$. Define $k_0 = \max\{k_1, k_2\}$. It is easy to see that $k_0 = k_1 \approx 1.62 \dots$.

4. PROOF OF THEOREM 1.1

From Theorem 2.6, we have

$$(4.1) \quad 0 < I_\beta(u_\beta) = \frac{1}{2} \int_{\mathbb{R}} |u''_\beta|^2 dx - \frac{\beta^2}{2} \int_{\mathbb{R}} |u'_\beta|^2 dx + \int_{\mathbb{R}} (e^{u_\beta} - u_\beta - 1) dx < C$$

where $C > 0$ is independent of β .

Let $\beta \in (0, \sqrt{2})$ be fixed. By Theorem 2.6, there exists a sequence $\beta_n \rightarrow \beta$ and a sequence of solutions of (1.4), called u_{β_n} , with Morse index at most one and the bound (4.1). Our main idea is to show that the limit of u_{β_n} exists and has uniform H^2 bound.

We will drop the subscript β for the sake of convenience.

By simple computations, it is easy to check that the function $\frac{e^u - u - 1}{u^2}$ is increasing if $u < 0$. Thus there is a unique $u_\star < 0$ such that

$$(4.2) \quad \frac{e^{u_\star} - u_\star - 1}{u_\star^2} = \frac{\beta^4 k_0^2}{8}.$$

Since $e^u - 1 - u \geq \frac{u^2}{2}$ for $u > 0$, we deduce that

$$(4.3) \quad \frac{e^u - u - 1}{u^2} \geq \frac{\beta^4 k_0^2}{8} \text{ for } u \geq u_\star$$

if

$$(4.4) \quad \beta \leq \frac{\sqrt{2}}{\sqrt{k_0}}.$$

Also we have

$$(4.5) \quad e^{u_*} \geq 1 + u_* + \frac{u_*^2}{2} e^{u_*}.$$

This implies that

$$e^{u_*} \leq \frac{\beta^4 k_0^2}{4}.$$

Our main idea is to bound the energy on the level sets $\{u \geq u_*\}$ and $\{u \leq u_*\}$. On the set $\{u \geq u_*\}$, we use the key inequality (3.2). On the set $\{u \leq u_*\}$, we will use Morse index information obtained in Theorem 2.6.

First, as a result of Remark 3.3 and the key inequalities (3.2) and (3.18) we have

$$(4.6) \quad \begin{aligned} & \frac{1}{2} \int_A |u''|^2 dx - \frac{\beta^2}{2} \int_A |u'|^2 dx + \int_A (e^u - u - 1) dx \\ & \geq \frac{1}{2} \int_A |u''|^2 dx - \frac{\beta^2}{2} \int_A |u'|^2 dx + \frac{\beta^4 k_0^2}{4} \int_A \frac{u^2}{2} dx \geq 0 \end{aligned}$$

where $A = \{u \geq u_*\}$. (Note that since u is a homoclinic, $A = (-\infty, b_0) \cup \cup_{j=1}^l (a_j, b_j) \cup (a_{l+1}, +\infty)$.)

Our main objective is then to show that in the complement of $A^c = \{u \leq u_*\}$,

$$(4.7) \quad \frac{1}{2} \int_{A^c} |u''|^2 dx - \frac{\beta^2}{2} \int_{A^c} |u'|^2 dx + \int_{A^c} (e^u - u - 1) dx \geq 0.$$

Let $A^c = \{u \leq u_*\} = \cup_{j=1}^m (a_j, b_j)$ where m is finite since u is homoclinic. Since Morse index of u is at most one, then except at most one interval (a_i, b_i) we must have, for $j \neq i$

$$(4.8) \quad \int_{a_j}^{b_j} |\varphi''|^2 dx - \beta^2 \int_{a_j}^{b_j} |\varphi'|^2 dx + \int_{a_j}^{b_j} e^u \varphi^2 dx \geq 0 \quad \forall \varphi \in C_0^2(a_j, b_j).$$

Without loss of generality let (4.8) hold in some interval (a, b) .

As e^u is an increasing function, in A^c , we have

$$(4.9) \quad e^u \leq e^{u_*}.$$

Note that from (4.8) we have

$$(4.10) \quad \int_a^b |\varphi''|^2 dx - \beta^2 \int_a^b |\varphi'|^2 dx + e^{u_*} \int_a^b \varphi^2 dx \geq 0 \quad \forall \varphi \in C_0^2(a, b).$$

(4.10) implies that the length of the interval $(b - a)$ can be controlled. In fact, we will have

$$(4.11) \quad \frac{b - a}{2} \leq a_*$$

where a_* depends on β .

On the other hand, if $v = u - u_*$, then we have

$$(4.12) \quad v'''' + \beta^2 v'' + e^{v+u_*} - 1 = 0.$$

Multiplying by v and integrating (4.12) we obtain

$$(4.13) \quad -v'v''|_a^b + \int_a^b (v'')^2 - \beta^2 \int_a^b (v')^2 + \int_a^b (e^{u_*+v} - 1)v = 0.$$

Integrating (2.4) we have

$$(4.14) \quad v'v''|_a^b - \frac{3}{2} \int_a^b (v'')^2 + \frac{\beta^2}{2} \int_a^b (v')^2 + \int_a^b (e^{u_*+v} - 1 - u_* - v) = 0.$$

Adding (4.13) and (4.14) we obtain that

$$(4.15) \quad \int_a^b (v'')^2 + \beta^2 \int_a^b (v')^2 = 2 \int_a^b (e^{u_*+v} - 1)v + 2 \int_a^b (e^{u_*+v} - u_* - v - 1)dx.$$

Substituting (4.15) into the energy over (a, b) we have

$$(4.16) \quad \begin{aligned} I_\beta|_{(a,b)}(u) &= \frac{1}{2} \int_a^b (u'')^2 - \frac{\beta^2}{2} \int_a^b (u')^2 + \int_a^b (e^u - u - 1)dx \\ &= \int_a^b (v'')^2 - \int_a^b (e^{u_*+v} - 1)v. \end{aligned}$$

Now we claim that

$$I_\beta|_{(a,b)}(u) \geq 0.$$

We argue by contradiction. If not, then $I_\beta|_{(a,b)}(u) \leq 0$. We have

$$(4.17) \quad \int_a^b (v'')^2 \leq \int_a^b (e^{u_*+v} - 1)v \leq \int_a^b (-v).$$

Without loss of generality we can consider (a, b) to be $(-a, a)$. Let

$$A = \int_{-a}^a (e^{u_*+v} - 1)v, \quad B = \int_{-a}^a (e^{u_*+v} - 1 - u_* - v), \quad \sigma := \frac{A}{B}$$

Then from (4.15) we have

$$(4.18) \quad \int_{-a}^a (u'')^2 + \beta^2 \int_{-a}^a (u')^2 \leq 2(1 + \sigma) \int_{-a}^a (e^{v+u_*} - u_* - v - 1).$$

Thus we have from (4.16)

$$(4.19) \quad \begin{aligned} I_\beta|_{(-a,a)}(u) &\geq \frac{1}{2} \left\{ \left(1 + \frac{1}{1+\sigma}\right) \int_{-a}^a (u'')^2 - \beta^2 \left(1 - \frac{1}{1+\sigma}\right) \int_{-a}^a (u')^2 \right\} \\ &\geq \frac{1}{2} \frac{1}{(1+\sigma)} \left\{ (2+\sigma) \int_{-a}^a (u'')^2 - \sigma \beta^2 \int_{-a}^a (u')^2 \right\} \\ &\geq c_0 \left\{ \int_{-a}^a (u'')^2 - \frac{\sigma}{2+\sigma} \beta^2 \int_{-a}^a (u')^2 \right\}. \end{aligned}$$

As a consequence, if $I_\beta|_{(-a,a)}(u) \leq 0$, then from (4.19) we obtain that

$$(4.20) \quad \int_{-a}^a (u'')^2 \leq \frac{\sigma}{2+\sigma} \beta^2 \int_{-a}^a (u')^2 dx.$$

Consider the following eigenvalue problem

$$(4.21) \quad \begin{cases} u'''' + \lambda^2 u'' = 0 & \text{in } (-a, a) \\ u = u'' = 0 & \text{on } \partial(-a, a). \end{cases}$$

By (4.20) the first eigenvalue λ_1^2 of (4.21) satisfies $\lambda_1^2 \leq \frac{\sigma\beta^2}{2+\sigma}$. But note that $u = A \cos(\lambda_1 x)$, with $\cos \lambda_1 a = 0$ implies that $\lambda_1 a = \frac{\pi}{2}$. This implies that $\lambda_1^2 a^2 = \frac{\pi^2}{4}$ and hence $\frac{\pi^2}{4} \leq \frac{\sigma}{2+\sigma} \beta^2 a^2$. As a result we obtain that

$$(4.22) \quad \beta a \geq \frac{\pi}{2} \sqrt{1 + \frac{2}{\sigma}}.$$

To estimate σ , let us notice that

$$(4.23) \quad \inf_{H^2(I) \cap H_0^1(I)} \frac{\int_{-a}^a (v'')^2}{(\int_{-a}^a v)^2} = \frac{15}{4a^5}.$$

In order to prove (4.23) consider the problem

$$(4.24) \quad \begin{cases} u'''' = 1 & \text{in } (-a, a) \\ u = u'' = 0 & \text{on } \partial(-a, a). \end{cases}$$

Then

$$u(x) = \frac{1}{24}(x^4 - a^4) - \frac{1}{4}a^2(x^2 - a^2)$$

and as a result we have

$$(4.25) \quad \int_{-a}^a u dx = \frac{1}{24}(6 + \frac{2}{5})a^5 = \frac{4}{15}a^5$$

and

$$(4.26) \quad \int_{-a}^a (u'')^2 dx = \frac{1}{24}(6 + \frac{2}{5})a^5 = \frac{4}{15}a^5.$$

Hence we have

$$(4.27) \quad \int_{-a}^a (v'')^2 dx \geq \frac{15}{4a^5} \left(\int_{-a}^a |v| \right)^2.$$

But from (4.17) we have

$$(4.28) \quad \int_{-a}^a (v'')^2 dx \leq \int_{-a}^a (-v) \leq \int_{-a}^a |v| dx.$$

This implies that

$$(4.29) \quad A \leq \int_{-a}^a |v| dx \leq \frac{4}{15}a^5.$$

But

$$B \geq A + (-1 - u_*)2a$$

and hence from (4.11) we have,

$$(4.30) \quad \frac{1}{\sigma} \geq 1 + \frac{15(-1 - u_*)}{2a_*^4}.$$

As a result of (4.22) we have

$$(4.31) \quad \frac{\sqrt{\frac{15(-1 - u_*)}{a_*^4} + 3}}{2\beta} \pi \leq a_*.$$

So as long as

$$(4.32) \quad \frac{\sqrt{\frac{15(-1-u_*)}{a_*^4} + 3}}{2} \pi > \beta a_*$$

holds we have a contradiction with (4.11).

Next we show that condition (4.32) holds when β is small. In fact, we have from (4.2) that for small β ,

$$(4.33) \quad u_* \approx -\frac{8}{\beta^4 k_0^2}$$

and hence $e^{u_*} \approx e^{-\frac{8}{\beta^4 k_0^2}} \approx 0$, we may assume that $e^{u_*} = 0$. Hence we solve the eigenvalue problem

$$(4.34) \quad \begin{cases} \varphi'''' + \beta^2 \varphi'' = 0 & \text{in } (-a, a) \\ \varphi = \varphi' = 0 & \text{on } \partial(-a, a). \end{cases}$$

Any even eigenfunction of (4.34) is $1 + \cos \beta x$ provided that $\beta a = \pi$. Hence from (4.10) and (4.34) we obtain

$$(4.35) \quad a \leq a_* = \frac{\pi}{\beta} + O(\beta)$$

and since $u_* \approx -\frac{8}{\beta^4 k_0^2}$, we obtain

$$(4.36) \quad 3 + \frac{15(-1-u_*)}{a_*^4} \geq 3 + \frac{120}{k_0^2 \pi^4} + O(\beta) \geq \frac{9}{2} + O(\beta)$$

which implies that (4.32) holds for β small.

We have thus proved that in A^c , except one interval,

$$(4.37) \quad I_\beta|_{(a,b)} \geq 0.$$

Let (a, b) be the exceptional interval in A^c . Then we have

$$(4.38) \quad \beta(b-a) < 4\pi.$$

In fact, if $\beta(b-a) \geq 4\pi$, then we can construct ψ_1 and ψ_2 having disjoint support such that

$$\begin{cases} \psi_1(x) = \cos \beta x & \text{in } (-\frac{\pi}{2\beta}, \frac{\pi}{2\beta}) \\ \psi_1(x) = \psi_1''(x) = 0 & \text{on } \partial(-\frac{\pi}{2\beta}, \frac{\pi}{2\beta}), \\ \psi_2(x) = \cos \beta x & \text{in } (\frac{\pi}{2\beta}, \frac{3\pi}{2\beta}) \\ \psi_2(x) = \psi_2''(x) = 0 & \text{on } \partial(\frac{\pi}{2\beta}, \frac{3\pi}{2\beta}) \end{cases}$$

and ψ_1 and ψ_2 contribute two to the Morse index of u , a contradiction to Theorem 2.6.

From (4.15), we have

$$(4.39) \quad \int_a^b (v'')^2 + \beta^2 (v')^2 \leq C + C \int_a^b |v|$$

which yields

$$(4.40) \quad |u| \leq |v| + |u_*| \leq C \text{ in } (a, b).$$

Then from (4.1) we have

$$(4.41) \quad -C \leq I_\beta|_{(a,b)}(u) \leq C.$$

Let $A' = A \setminus (a, b)$. Then

$$(4.42) \quad 0 < \int_{A'} (u'')^2 - \beta^2 \int_{A'} (u')^2 + \int_{A'} (e^u - u - 1) dx \leq C$$

and this implies that

$$(4.43) \quad \int_{A'} (e^u - u - 1) dx + \int_{A'} ((u'')^2 + (u')^2 + u^2) \leq C$$

and hence $|u| \leq C$ in A . Multiplying (1.4) by u and integrating we obtain

$$(4.44) \quad \int_{\mathbb{R}} (u'')^2 - \beta^2 \int_{\mathbb{R}} (u')^2 + \int_{\mathbb{R}} (e^u - 1)u = 0.$$

From (4.1) and (4.44) we have

$$(4.45) \quad -\frac{1}{2} \int_{\mathbb{R}} (e^u - 1)u + \int_{\mathbb{R}} (e^u - u - 1) dx < C$$

and this implies

$$(4.46) \quad \int_{u < 0} [(e^u - u - 1) - \frac{1}{2}(e^u - 1)u] < C.$$

Moreover, using (4.15) we obtain

$$(4.47) \quad \int_{\mathbb{R}} (u'')^2 + \beta^2 \int_{\mathbb{R}} (u')^2 = 2 \int_{\mathbb{R}} (e^u - 1)u + 2 \int_{\mathbb{R}} (e^u - u - 1) \leq C.$$

This implies

$$(4.48) \quad \|u_\beta\|_{H^2(\mathbb{R})} \leq C.$$

Let $\beta \in (0, \sqrt{2})$ such that there exists $\beta_n \rightarrow \beta$ as $n \rightarrow \infty$ and for $\beta = \beta_n$ (1.4) has a solution. Hence we have

$$u_{\beta_n}'''' + \beta_n^2 u_{\beta_n}'' + e^{u_{\beta_n}} - 1 = 0.$$

Also we have $\|u_{\beta_n}\|_{H^2(\mathbb{R})} \leq C$ and hence $u_{\beta_n} \rightharpoonup u_\beta$ in $H^2(\mathbb{R})$ and as a result, we have $u_{\beta_n} \rightarrow u_\beta$ in $L_{loc}^p(\mathbb{R})$ as $n \rightarrow +\infty$ for all p . In particular, $u_{\beta_n} \rightarrow u_\beta$ in $C_{loc}^1(\mathbb{R})$. Hence $u_{\beta_n}(x) \rightarrow u_\beta(x)$ pointwise almost everywhere. Thus $e^{u_{\beta_n}(x)} \rightarrow e^{u_\beta(x)}$ almost everywhere. As $n \rightarrow +\infty$, we have

$$u_\beta'''' + \beta^2 u_\beta'' + e^{u_\beta} - 1 = 0.$$

Finally we prove that u_β is nontrivial, i.e., $u_\beta \not\equiv 0$. As u_{β_n} is a solution of (1.4) from (4.44) we have

$$(4.49) \quad \int_{\mathbb{R}} (u_{\beta_n}'')^2 - \beta_n^2 \int_{\mathbb{R}} (u_{\beta_n}')^2 = - \int_{\mathbb{R}} (e^{u_{\beta_n}} - 1)u_{\beta_n} dx.$$

Invoking Fourier transform technique as in Lemma 2.2 we have

$$(4.50) \quad \int_{\mathbb{R}} (u_{\beta_n}'')^2 - \beta_n^2 \int_{\mathbb{R}} (u_{\beta_n}')^2 \geq -\frac{\beta_n^4}{4} \int_{\mathbb{R}} u_{\beta_n}^2.$$

Hence we have

$$(4.51) \quad -\frac{\beta_n^4}{4} \int_{\mathbb{R}} u_{\beta_n}^2 \leq - \int_{\mathbb{R}} (e^{u_{\beta_n}} - 1)u_{\beta_n} dx.$$

This implies

$$(4.52) \quad \int_{\mathbb{R}} \left[\frac{\beta_n^4}{4} u_{\beta_n}^2 - (e^{u_{\beta_n}} - 1) u_{\beta_n} \right] dx \geq 0.$$

As a result, there exists $x_n \in \mathbb{R}$ such that

$$\frac{\beta_n^4}{4} u_{\beta_n}^2(x_n) \geq (e^{u_{\beta_n}(x_n)} - 1) u_{\beta_n}(x_n).$$

Note that this can only happen when $u_{\beta_n}(x_n) < 0$. Hence

$$\frac{\beta_n^4}{4} \geq \frac{e^{u_{\beta_n}(x_n)} - 1}{u_{\beta_n}(x_n)} \geq e^{u_{\beta_n}(x_n)}.$$

Thus $e^{u_{\beta_n}(x_n)} \leq \frac{\beta_n^4}{4}$ and hence $u_{\beta_n}(x_n) \leq \ln \frac{\beta_n^4}{4}$. Hence there exists $x_0 \in \mathbb{R}$ such that $u_{\beta}(x_0) \leq \ln \frac{\beta^4}{4} < 0$. This implies u_{β} is a nontrivial solution of (1.4).

5. RANGE OF β AND DECAY ESTIMATES

In this section, we first find explicit bound for β so that (4.32) holds, and then we prove the decay estimate.

5.1. Estimate of β . First, we find a_* . We recall the following eigenvalue problem

$$(5.1) \quad \begin{cases} \varphi'''' + \beta^2 \varphi'' + e^{u_*} \varphi = 0 & \text{in } (-a, a) \\ \varphi(\pm a) = \varphi'(\pm a) = 0. \end{cases}$$

Any even solution of (5.1) can be written as $u(x) = A \cos \mu_1 x + B \cos \mu_2 x$ where

$$\mu_1 = \sqrt{\frac{\beta^2}{2} - \sqrt{\frac{\beta^4}{4} - e^{u_*}}}$$

and

$$\mu_2 = \sqrt{\frac{\beta^2}{2} + \sqrt{\frac{\beta^4}{4} - e^{u_*}}}.$$

Then they must satisfy

$$(5.2) \quad \mu_2 \tan \mu_2 a = \mu_1 \tan \mu_1 a.$$

Since $\mu_1 < \mu_2$, the function $\mu \tan \mu a$ is increasing where $a \in (0, \frac{\pi}{2\mu_2})$. Hence (5.2) admits a solution in $(\frac{\pi}{2\mu_2}, \frac{3\pi}{2\mu_2})$. When $e^{u_*} \ll 1$, μ_1 is close to zero, then

$$\mu_2 \tan \mu_2 a \approx \mu_1^2 a.$$

Let $\mu_2 a = \pi + t$. Then from (5.2)

$$t \leq \tan t = \frac{\mu_1}{\mu_2} \tan \frac{\mu_1}{\mu_2} (\pi + t), \quad t \in (0, \frac{\pi}{2}).$$

Thus we obtain that $a \leq a_*$ where

$$(5.3) \quad a_* := \frac{\pi}{\mu_2} + \frac{\mu_1}{\mu_2} \tan \frac{3\mu_1}{2\mu_2} \pi.$$

The condition (4.32) can be checked numerically using (4.2) to find an approximate bound for u_* and we find that the numerical bound for (4.32) to hold if $\beta \leq \beta_* \approx 0.742 \dots$.

5.2. Decay estimates of (1.4). Note that $u(x) \rightarrow 0$ as $x \rightarrow \pm\infty$. Hence

$$(5.4) \quad \frac{e^u - 1}{u} \rightarrow 1$$

Hence the limiting equation for fixed β at infinity is given by

$$(5.5) \quad w'''' + \beta^2 w'' + w = 0.$$

Note that this is a linear problem and the roots of the

$$(5.6) \quad m^4 + \beta^2 m^2 + 1 = 0$$

and hence

$$(5.7) \quad \left(m^2 + \frac{\beta^2}{2}\right)^2 = \left(1 - \frac{\beta^4}{4}\right)i^2$$

where $i = \sqrt{-1}$. Define $n = m^2$ then we have

$$(5.8) \quad n = -\frac{\beta^2}{2} \pm \left(1 - \frac{\beta^4}{4}\right)i^2.$$

Define $\cos 2\eta = -\frac{\beta^2}{2}$. Then we can write (5.8) as

$$n = (\cos 2\eta + i \sin 2\eta) = (\cos \eta + i \sin \eta)^2$$

where $\eta \in (\frac{\pi}{4}, \frac{\pi}{2})$ when $x \rightarrow -\infty$ and $\eta \in (-\frac{\pi}{2}, -\frac{\pi}{4})$ when $x \rightarrow +\infty$ as we are looking for decaying solutions. This implies that the four roots of (5.7) are precisely $m = e^{\pm i\eta}$ and $\bar{m} = e^{\pm i\bar{\eta}}$. If

$$m = e^{i\eta} = \tau + i\delta$$

where $\tau = \cos \eta$ and $\delta = \sin \eta$. Hence the general solution of (5.5) decaying at $\pm\infty$ is given by

$$(5.9) \quad w(x) = e^{\tau x} \cos(\delta x + b) \chi_{\{x < 0\}} + e^{-\tau x} \cos(\delta x + d) \chi_{\{x > 0\}}$$

for some $b, d \in \mathbb{R}$, χ denotes the characteristic function. Also note that τ depends on β . As a result, u decays exponentially for each $\beta > 0$.

6. PROOF OF THEOREM 1.2

The ideas used in proving Theorem 1.1 can be readily extended to (1.7). We make a change of variable $u - 1$ in (1.7). Then the equation transforms into

$$(6.1) \quad \begin{cases} u'''' + \beta^2 u'' + u^3 + 3u^2 + 2u = 0 & \text{in } \mathbb{R} \\ u(x) > -2 & \text{in } \mathbb{R} \\ u \in H^2(\mathbb{R}). \end{cases}$$

Define $J_\beta : H^2(\mathbb{R}) \rightarrow \mathbb{R}$ as

$$J_\beta(u) = \frac{1}{2} \int_{\mathbb{R}} (u'')^2 - \frac{\beta^2}{2} \int_{\mathbb{R}} (u')^2 + \frac{1}{4} \int_{\mathbb{R}} u^4 + \int_{\mathbb{R}} u^3 + \int_{\mathbb{R}} u^2.$$

Smets-van den Berg [29] proved that for *almost all* $\beta \in (0, \sqrt{8})$, problem (6.1) has a homoclinic solution with

$$(6.2) \quad u > -2.$$

Let $u_\star$ be such that

$$(6.3) \quad u_\star = -2 + \frac{k_0}{\sqrt{2}} \beta^2$$

By our assumption $\beta^2 < \frac{\sqrt{2}}{k_0}$, we have $u_* \leq -1$. In $A = \{u \geq u_*\}$, we have

$$(6.4) \quad I_A \geq \frac{1}{2} \int_A (u'')^2 - \frac{\beta^2}{2} \int_{A_2} (u')^2 + \frac{k_0^2 \beta^4}{8} \int_A u^2 \geq 0$$

by (3.2) and (3.18).

Let $A^c = \{u \leq u_*\} = \cup_{j=1}^k (a_j, b_j)$, k is finite since u is homoclinic. Let (a, b) be one of the intervals in A^c . If $v = u - u_*$, then we have

$$(6.5) \quad v'''' + \beta^2 v'' + u(u+1)(u+2) = 0.$$

Multiplying by v and integrating (6.5) we obtain

$$(6.6) \quad -v'v'' \Big|_a^b + \int_a^b (v'')^2 - \beta^2 \int_a^b (v')^2 + \int_a^b u(u+1)(u+2)(u-u_*) = 0.$$

Similar to (2.4), we obtain a pointwise identity

$$(6.7) \quad u'(x)u'''(x) - \frac{(u''(x))^2}{2} + \frac{\beta^2}{2}(u'(x))^2 + \frac{1}{4}u^2(x)(u(x)+2)^2 = 0.$$

Integrating (2.4) we have

$$(6.8) \quad v'v'' \Big|_a^b - \frac{3}{2} \int_a^b (v'')^2 + \frac{\beta^2}{2} \int_a^b (v')^2 + \frac{1}{4} \int_a^b u^2(u+2)^2 = 0.$$

Adding (6.6) and (6.8) we obtain that

$$(6.9) \quad \int_a^b (v'')^2 + \beta^2 (v')^2 = 2 \int_a^b u(u+1)(u+2)(u-u_*) + \frac{1}{2} \int_a^b u^2(u+2)^2 dx.$$

Substituting (6.9) into the energy over (a, b) we have

$$(6.10) \quad \begin{aligned} I_\beta \Big|_{(a,b)} (u) &= \frac{1}{2} \int_a^b (u'')^2 - \frac{\beta^2}{2} \int_a^b (u')^2 + \int_a^b \frac{1}{4} u^2 (u+2)^2 dx \\ &= \int_a^b (v'')^2 - \int_a^b u(u+1)(u+2)(u-u_*). \end{aligned}$$

Now we claim that

$$I_\beta \Big|_{(a,b)} (u) \geq 0.$$

In fact, we have $u+2 > 0, u < 0, u+1 \leq u_*+1 \leq 0$, and hence

$$(6.11) \quad u(u+1)(u+2)(u-u_*) \leq 0 \text{ on } (a, b)$$

This implies $I_\beta \Big|_{(a,b)} (u) \geq 0$ and hence

$$(6.12) \quad I_\beta \Big|_A (u) \geq 0.$$

The rest of the proof is similar to that of Theorem 1.1. We omit the details. $\square$

ACKNOWLEDGEMENT

The research was partially supported by an ARC grant, an Earmarked grant from RGC of Hong Kong and a Focused Research Scheme from CUHK.

REFERENCES

- [1] G. ADAMS, L. LIN; Beam on a tensionless elastic foundation. *Journal of Engineering Mechanics*. 113 (1986), no. 2, 542–553.
- [2] N. BOCCARA, D. LANANGÉVIN, J. MEUNIER; Physics of Amphiphilic layer. *Springer Proc. of Physics* Vol. 21 (1987).
- [3] D. BONHEURE; Multitransition kinks and pulses for fourth order equations with a bistable nonlinearity. *Ann. Inst. H. Poincaré Anal. Non Linéaire* 21 (2004), no. 3, 319–340.
- [4] D. BONHEURE, D. SANCHEZ, M. TARALLO, S. TERRACINI; Heteroclinic connections between nonconsecutive equilibria of a fourth order differential equation. *Calc. Var. Partial Differential Equations* 17 (2003), no. 4, 341–356.
- [5] B. BREUER, J. HORÁK, P.J. MCKENNA, M. PLUM; A computer-assisted existence and multiplicity proof for travelling waves in a nonlinearly supported beam. *J. Differential Equations* 224 (2006), no. 1, 60–97.
- [6] B. BUFFONI; Periodic and homoclinic orbits for Lorentz-Lagrangian systems via variational methods. *Nonlinear Anal.* 26 (1996), no. 3, 443–462.
- [7] B. BUFFONI, A. CHAMPNEYS, J. TOLAND; Bifurcation and coalescence of a plethora of homoclinic orbits for a Hamiltonian system. *J. Dynam. Differential Equations* 8 (1996), no. 2, 221–279.
- [8] A. BLACKMORE, G. HUNT; Principle of localised buckling for a strut on an elastoplastic foundation. *Trans. of AMSE.* 63 (1996), no. 2, 234–239.
- [9] A. CHAMPNEYS, P. J. MCKENNA; On solitary waves of a piecewise linear suspended beam model. *Nonlinearity* 10 (1997), no. 6, 1763–1782.
- [10] A. CHAMPNEYS, P. J. MCKENNA, P. ZEGELING; Solitary waves in nonlinear beam equations: stability, fission and fusion. The theme of solitary waves and localization phenomena in elastic structures. *Nonlinear Dynam.* 21 (2000), no. 1, 31–53.
- [11] Y. CHEN, P. J. MCKENNA; Travelling waves in a nonlinearly suspended beam: theoretical results and numerical observations. *J. Diff. Equations* 136 (1997), no. 2, 325–355.
- [12] Y. CHEN, P.J. MCKENNA; Travelling waves in a nonlinearly suspended beam: some computational results and four open questions. *Philos. Trans. Roy. Soc. London Ser. A* 355 (1997), no. 1732, 2175–2184.
- [13] E. DANCER; Finite Morse index solutions of supercritical problems, *J. Reine Angew. Math.* 620(2008), 213–233.
- [14] R. DEVANEY; Homoclinic orbits in Hamiltonian systems. *J. Differential Equations* 21 (1976), no. 2, 431–438.
- [15] I. EKELAND, N. GHOUSSEUB; Selected new aspects of the calculus of variations in the large. *Bull. Amer. Math. Soc. (N.S.)* 39 (2002), no. 2, 207–265.
- [16] A. FARINA, On the classification of solutions of Lane-Emden equation on unbounded domains of $\mathbb{R}^N$, *J. Math. Pures Appl.* 87(2007), no.5, 537–561.
- [17] N. GHOUSSEUB; Duality and perturbation methods in critical point theory. *Cambridge tracts in Mathematics* (1993) .
- [18] G. GOMPPER, M. SCHICK; Self-assembling amphiphilic systems *PTCP* Vol.16 AP, 1994.
- [19] A. LAZER, P.J. MCKENNA; Large-amplitude periodic oscillations in suspension bridges: some new connections with nonlinear analysis. *SIAM Rev.* 32 (1990), no. 4, 537–578.
- [20] J. LEGA, J. MOLONEY, A. NEWELL; Swift-Hohenberg for lasers. *Physics Review Letters* Volume 73, Issue 4, (1994) p. 2978–2981.
- [21] P.J. MCKENNA; Large-amplitude periodic oscillations in simple and complex mechanical systems: outgrowths from nonlinear analysis. *Milan J. Math.* 74(2006), 79–115.
- [22] P. J. MCKENNA, W. WALTER; Travelling waves in a suspension bridge. *SIAM J. Appl. Math.* 50 (1990), no. 3, 703–715.
- [23] Y. POMEAU, P. MANNEVILLE; Wavelength selection in cellular flows. *Physics Letters A* Volume 75, Issue 4, (1980) p. 296–298.
- [24] L.A. PELETIER, W.C. TROY; Spatial Patterns, Higher Order Models in Physics and Mechanics, Birkhauser, 2001.
- [25] L.A. PELETIER, W.C. TROY; A topological shooting method and the existence of kinks of the extended Fisher-Kolmogorov equation. *Topol. Methods Nonlinear Anal.* 6 (1995), no. 2, 331–355.

- [26] L.A. PELETIER, W.C. TROY; Spatial patterns described by the extended Fisher-Kolmogorov (EFK) equation: kinks. *Differential Integral Equations* 8 (1995), no. 6, 1279–1304.
- [27] L.A. PELETIER, W.C. TROY; Multibump periodic travelling waves in suspension bridges. *Proc. Roy. Soc. Edinburgh Sect. A* 128 (1998), no. 3, 631–659.
- [28] A. POLYAKOV; Gauge fields and strings. *Harwood, NY* (1988).
- [29] D. SMETS, J. B. VAN DEN BERG; Homoclinic solutions for Swift-Hohenberg and suspension bridge type equations. *J. Differential Equations* 184 (2002), no. 1, 78–96.;
- [30] M. Struwe, The existence of surfaces of constant mean curvature with free boundaries, *Acta Math.* 160(1988), 19-64.
- [31] J.B. SWIFT, P.C. HOHENBERG, Hydrodynamic fluctuations at the convective instability, *Phys. Rev. A* 51 (1977), 319-328.
- [32] J. B. van den Berg; The phase-plane picture for a class of fourth-order conservative differential equations. *J. Differential Equations* 161 (2000), no. 1, 110–153.

S. SANTRA, SCHOOL OF MATHEMATICS AND STATISTICS, THE UNIVERSITY OF SYDNEY, NSW 2006, AUSTRALIA.

E-mail address: `sanjiban@maths.usyd.edu.au`

J. WEI, DEPARTMENT OF MATHEMATICS, THE CHINESE UNIVERSITY OF HONG KONG, SHATIN, HONG KONG.

E-mail address: `wei@math.cuhk.edu.hk`